

NETYLIN

(Netilmicin)

0.3% w/v

Sterile Ophthalmic Solution

For Your Eyes Only



INNVOTEK

نیٹیلین
آئی ڈراپس

(نیٹیل مائیسین)

0.3% w/v

COMPOSITION:

Each mL contains:

Netilmicin sulphate BP equivalent to

Netilmicin..... 3mg.

(Innovator's Specs)

Inactives and purified water.

PHARMACOTHERAPEUTIC CLASS:

Netilmicin is an aminoglycoside antibiotic.

THERAPEUTIC INDICATIONS:

Topical treatment of external infections of the eye and its adnexa caused by germs sensitive to Netilmicin, including those producing adenylating or phosphorylating enzymes.

DOSAGE AND ADMINISTRATION:

Instill 1 to 2 drops in the conjunctival sac 3 times daily or as directed by the physician.

PRECAUTIONS FOR USE:

Prolonged use of topical products may determine sensitization phenomena. In such cases discontinue treatment and consult the physician to start an adequate therapy. In infants and pregnant women the product should be administered in case of real need and under strict medical control. The use of the product is not recommended during lactation. Netylin eye drops is not injectable: therefore, it must not be injected subconjunctivally or introduced in the anterior chamber.

PHARMACOLOGICAL INTERACTIONS:

No significant pharmacological interactions have ever been reported further to the use of Netilmicin as eye drops. Concomitant administration, even topical and especially intracavitary, of other potentially nephro and ototoxic antibiotics may increase the risk of such effects. An increase in potential nephrotoxicity of some aminoglycosidcs is reported further to subsequent or concomitant use, which should be avoided of other potentially nephrotoxic drugs such as cisplatin, polymyxin B, colistin, viomycin, streptomycin, vancomycin, other aminoglycosides and some cephalosporins (cephaloridin) or potent diuretics such as ethacrynic acid and furosemide for their effects on kidneys. In vitro, the association of an aminoglycoside with a betalactam antibiotic (penicillins or cephalosporins) may cause a reciprocal and relevant inactivation. A reduction of half-life or plasma levels of aminoglycoside occurred in patients suffering from renal insufficiency and in some patients with normal renal activity, even if any aminoglycoside and a penicillin like antibiotic were administered' by two different routes.

SPECIAL WARNINGS:

Prolonged use of topical antibiotics may determine over growth of resistant micro-organisms. Should no significant clinical improvement be reported within a relatively short period of time or should any irritation or sensitization phenomena occurs, is necessary to discontinue treatment and start an adequate therapy.

UNDESIRABLE EFFECTS:

Netylin may occasionally cause slight and transient ocular irritation or sensitization phenomena such as edema, urticarial eruptions and pruritus. Patients are kindly requested to inform their doctor or chemist for any undesirable effect not described in this leaflet.

CONTRA-INDICATIONS:

Patient's hypersensitivity to the drug components or to any other aminoglycoside.

INSTRUCTIONS:

Store between 8°C to 25°C. Protect from sunlight and heat. Keep out of the reach of children. Use the solution within one month after first opening the bottle. Use as directed by the physician.

PACK PRESENTATION:

Netylin 0.3% w/v sterile ophthalmic solution is available in 5mL plastic dropper bottle.

طریقہ استعمال: ڈاکٹر کی ہدایت کے مطابق استعمال کریں۔

ہدایات: دو اکوروشنی، گرمی اور نمی سے بچائیں۔ تمام ادویات بچوں کی پہنچ سے دور رکھیں۔

۸ سے ۲۵ ڈگری سینٹی گریڈ درجہ حرارت پر رکھیں۔

پہلی دفعہ بوتل کھولنے کے بعد دو کو ایک ماہ تک استعمال کیا جا سکتا ہے۔



INNVOTEK

Manufactured by:

Innovatek Pharmaceuticals

Plot # 35-Industrial Estate Triangle,
Kahuta Road, Islamabad-Pakistan.

ISO Certified Company